arXiv:1303.5110v2 [quant-ph] 8 Apr 2013

One-norm geometric quantum discord under decoherence

J. D. Montealegre,[1,*] F. M. Paula,[1,†] A. Saguia,[1,‡] and M. S. Sarandy[1,§]
[1]*Instituto de Física, Universidade Federal Fluminense,*
Av. Gal. Milton Tavares de Souza s/n, Gragoatá, 24210-346, Niterói, RJ, Brazil

(Dated: April 10, 2013)

Geometric quantum discord is a well-defined measure of quantum correlation if Schatten 1-norm (trace norm) is adopted as a distance measure. Here, we analytically investigate the dynamical behavior of the 1-norm geometric quantum discord under the effect of decoherence. By starting from arbitrary Bell-diagonal mixed states under Markovian local noise, we provide the decays of the quantum correlation as a function of the decoherence parameters. In particular, we show that the 1-norm geometric discord exhibits the possibility of double sudden changes and freezing behavior during its evolution. For nontrivial Bell-diagonal states under simple Markovian channels, these are new features that are in contrast with the Schatten 2-norm (Hilbert-Schmidt) geometric discord. The necessary and sufficient conditions for double sudden changes as well as their exact locations in terms of decoherence probabilities are provided. Moreover, we illustrate our results by investigating decoherence in quantum spin chains in the thermodynamic limit.

PACS numbers: 03.65.Ud, 03.67.Mn, 75.10.Jm

I. INTRODUCTION

Geometry plays a fundamental role in physics. In modern quantum theory, this idea has been perpetrated through many proposals of geometrization of quantum information resources, e.g., entanglement [1–4]. In recent years, geometric approaches have also been extended for more general quantum correlation measures. In this scenario, special focus has been given to quantum discord (QD) (see Refs. [5–7] for recent reviews), which is an information-theoretic measure of non-classical correlations originally proposed by Ollivier and Zurek [8] through an entropic (non-geometric) approach.

The geometrization of QD, named as geometric quantum discord (GQD), has been proposed by Dakić, Vedral, and Brukner [9], mainly motivated by the difficulty of extracting analytical solutions for the entropic version of QD. GQD, as originally defined, can be seen as a deformation of the entropic QD [10] and quantifies the amount of quantum correlations in terms of its minimal Hilbert-Schmidt distance from the set of classical states. Its evaluation is analytically computable for general two-qubit states [9] as well as for arbitrary bipartite states [11–13]. Moreover, it has been shown to exhibit operational significance in specific quantum communication protocols (see, e.g., Refs. [14, 15]). Despite those remarkable features, GQD is known to be sensitive to the choice of distance measures (see, e.g., Ref. [16]). In turn, as recently pointed out [17–19], GQD as proposed in Ref. [9] cannot be regarded as a good measure for the quantumness of correlations, since it may increase under local operations on the unmeasured subsystem. In particular, it has explicitly been shown by Piani in Ref. [19] that the simple introduction of a factorized local ancillary state on the unmeasured party changes the geometric discord by a factor given by the purity of the ancilla. From a technical point of view, the root of this drawback is the lack of contractivity of the Hilbert-Schmidt norm under trace-preserving quantum channels. Remarkably, this problem can be completely solved if Schatten 1-norm (trace norm) is adopted as a distance measure. Indeed, concerning general Schatten p-norms for a geometric definition of QD [20], it has been shown in Ref. [21] that the 1-norm is the only p-norm able to consistently define GQD (see also Ref. [22]).

In this work, we investigate the 1-norm GQD under the effect of decoherence. More specifically, we consider Bell-diagonal states going through Markovian local quantum channels that preserve the Bell-diagonal form. For these states, we analytically provide the 1-norm GQD as a function of the decoherence probability. In particular, we show that the 1-norm GQD may exhibit double sudden changes and freezing behavior during its evolution. Concerning double sudden changes, its occurrence for Bell-diagonal states under the simplest Markovian evolutions is a new feature that have not been found for both the entropic [23] and the Schatten 2-norm geometric [24] QD. Specifically about the freezing behavior for general Bell-diagonal states under Markovian noise, it also occurs for the entropic QD [36], being interpreted as a sudden transition between classical and quantum decoherence. We illustrate double sudden changes and freezing of the 1-norm GQD through simple examples. Moreover, we also consider the dynamical properties of quantum correlations in the XXZ quantum spin chain in the thermodynamic limit, showing a connection between the occurrence of sudden changes and the quantum phases of the XXZ model.

---

[*] jdmontealegrer@if.uff.br
[†] fagner@if.uff.br
[‡] amen@if.uff.br
[§] msarandy@if.uff.br

II. GQD AND SCHATTEN 1-NORM

Let us begin by considering a bipartite system AB in a Hilbert space $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B$. The system is characterized by quantum states described by density operators $\rho \in \mathcal{B}(\mathcal{H})$, where $\mathcal{B}(\mathcal{H})$ is the set of bound, positive-semidefinite operators acting on $\mathcal{H}$ with trace given by $\mathrm{Tr}\,[\rho] = 1$. The 1-norm GQD between A and B is defined through the trace distance between ρ and the closest classical-quantum state ρ_c, reading

$$D_G(\rho) = \min_{\Omega_0} \left\| \rho - \rho_c \right\|_1 , \tag{1}$$

where $\left\| X \right\|_1 = \mathrm{Tr}\left[\sqrt{X^\dagger X}\right]$ is the 1-norm (trace norm) and Ω_0 is the set of classical-quantum states, whose general form is given by

$$\rho_c = \sum_k p_k \Pi_k^A \otimes \rho_k^B , \tag{2}$$

with $0 \leq p_k \leq 1$ ($\sum_k p_k = 1$), $\{\Pi_k^a\}$ denoting a set of orthogonal projectors for subsystem A, and ρ_k^B a general reduced density operator for subsystem B.

We will focus here in the particular case of two-qubit Bell diagonal states, whose density operator presents the form

$$\rho = \frac{1}{4}\left[I \otimes I + \vec{c} \cdot (\vec{\sigma} \otimes \vec{\sigma}) \right] , \tag{3}$$

where I is the identity matrix, $\vec{c} = (c_1, c_2, c_3)$ is a three-dimensional vector such that $-1 \leq c_i \leq 1$ and $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ is a vector composed by Pauli matrices. In matrix notation, we have

$$\rho = \frac{1}{4} \begin{pmatrix} 1 + c_3 & 0 & 0 & c_1 - c_2 \\ 0 & 1 - c_3 & c_1 + c_2 & 0 \\ 0 & c_1 + c_2 & 1 - c_3 & 0 \\ c_1 - c_2 & 0 & 0 & 1 + c_3 \end{pmatrix} . \tag{4}$$

The minimization over the whole set of classical states was obtained for GQD as defined by the 2-norm [11] and by the relative entropy [25]. For those cases, it can be proved that the minimal state is a decohered (measured) state, i.e. the minimization runs over classical states of the form $\rho_c = \Phi(\rho)$, where

$$\Phi(\rho) = \sum_{k=-,+} \left(\Pi_k \otimes I \right) \rho \left(\Pi_k \otimes I \right) , \tag{5}$$

with

$$\Pi_\pm = \frac{1}{2}\left(I \pm \vec{n} \cdot \vec{\sigma} \right) \tag{6}$$

and $\vec{n} = (n_1, n_2, n_3)$ is an unitary vector. For the 1-norm GQD, minimization over decohered states has also been shown to be sufficient for the case of Bell-diagonal states [21]. In this case, the 1-norm GQD can be analytically computed, yielding [21]

$$D_G(\rho) = \mathrm{int}\left[\left| c_1 \right|, \left| c_2 \right|, \left| c_3 \right| \right] , \tag{7}$$

where $\mathrm{int}\left[\left| c_1 \right|, \left| c_2 \right|, \left| c_3 \right| \right]$ describes the *intermediate* result among the absolute values of the correlation functions c_1, c_2 and c_3. More specifically, for $|c_i| \leq |c_j| \leq |c_k|$ ($i, j, k \in \{1, 2, 3\}$), we attribute $|c_j| = \mathrm{int}\left[|c_i|, |c_j|, |c_k| \right]$. Remarkably, this is equivalent to the *negativity of quantumness*, which is a measure of nonclassicality recently introduced in Refs. [26, 27] and experimentally discussed in Ref. [28]. Then, for Bell-diagonal states, by investigating the general behavior of the 1-norm GQD under decoherence, we are automatically providing the behavior of the negativity of quantumness under decoherence.

III. 1-NORM GQD UNDER DECOHERENCE

We will consider the system-environment interaction through the operator-sum representation formalism [29]. In this context, we will take the evolution of a quantum state ρ as described by a trace-preserving quantum operation $\varepsilon(\rho)$, which is given by

$$\varepsilon(\rho) = \sum_{i,j} \left(E_i \otimes E_j \right) \rho \left(E_i \otimes E_j \right)^\dagger , \tag{8}$$

where $\{E_k\}$ is the set of Kraus operators associated to a decohering process of a single qubit, with the trace-preserving condition reading $\sum_k E_k^\dagger E_k = I$. We provide in Table I a list of Kraus operators for a variety of channels considered in this work.

	Kraus operators
BF	$E_0 = \sqrt{1 - p/2}\, I, E_1 = \sqrt{p/2}\, \sigma_1$
PF	$E_0 = \sqrt{1 - p/2}\, I, E_1 = \sqrt{p/2}\, \sigma_3$
BPF	$E_0 = \sqrt{1 - p/2}\, I, E_1 = \sqrt{p/2}\, \sigma_2$
GAD	$E_0 = \sqrt{p} \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{1-\gamma} \end{pmatrix}, E_2 = \sqrt{1-p} \begin{pmatrix} \sqrt{1-\gamma} & 0 \\ 0 & 1 \end{pmatrix}$ $E_1 = \sqrt{p} \begin{pmatrix} 0 & \sqrt{\gamma} \\ 0 & 0 \end{pmatrix}, E_3 = \sqrt{1-p} \begin{pmatrix} 0 & 0 \\ \sqrt{\gamma} & 0 \end{pmatrix}$

TABLE I. Kraus operators for the quantum channels: bit flip (BF), phase flip (PF), bit-phase flip (BPF), and generalized amplitude damping (GAD), where p and γ are decoherence probabilities.

The decoherence processes BF, PF, and BPF in Table I preserve the Bell-diagonal form of the density operator ρ. For the case of GAD, the Bell-diagonal form is kept for arbitrary γ and $p = 1/2$. In this situation, we can write the quantum operation $\varepsilon(\rho)$ as

$$\varepsilon(\rho) = \frac{1}{4} \begin{pmatrix} 1 + c_3' & 0 & 0 & c_1' - c_2' \\ 0 & 1 - c_3' & c_1' + c_2' & 0 \\ 0 & c_1' + c_2' & 1 - c_3' & 0 \\ c_1' - c_2' & 0 & 0 & 1 + c_3' \end{pmatrix} , \tag{9}$$

where the values of the correlation vector $\vec{c'} = (c'_1,\ c'_2,\ c'_3)$ are given in Table II.

Channel	c'_1	c'_2	c'_3
BF	c_1	$c_2(1-p)^2$	$c_3(1-p)^2$
PF	$c_1(1-p)^2$	$c_2(1-p)^2$	c_3
BPF	$c_1(1-p)^2$	c_2	$c_3(1-p)^2$
GAD	$c_1(1-\gamma)$	$c_2(1-\gamma)$	$c_3(1-\gamma)^2$

TABLE II. Correlation functions for the quantum operations: bit flip (BF), phase flip (PF), bit-phase flip (BPF), and generalized amplitude damping (GAD). For GAD, we fixed $p = 1/2$.

Since the Bell-diagonal form is preserved in Eq. (9), we can directly obtain the 1-norm GQD from Table II by using that

$$D_G[\varepsilon(\rho)] = \text{int}\left[|c'_1|\,,|c'_2|\,,|c'_3|\right]. \qquad (10)$$

Then, we can obtain a general pattern of the 1-norm GQD decay as a function of the decoherence parameters. In particular, depending on both the original density operator ρ and the quantum channel, we may have sudden changes in the decay rate of $D_G(\rho)$. This will occur whether there are crossings among the components $|c'_1|$, $|c'_2|$, and $|c'_3|$ as a function of p or γ. Moreover, additional effects may appear, which corresponds to *double sudden changes* and *freezing behavior*. More specifically, we can identify the following general behavior for the dynamics in Table II:

(i) BF, PF, and BPF: Double sudden changes occur if and only if the constant component $|c_k|$ for the channel is nonvanishing and is the lowest absolute value of the components of the correlation vector $\vec{c}$, with the two other components exhibiting distinct values ($|c_k| = |c_1|$ for BF, $|c_k| = |c_3|$ for PF, and $|c_k| = |c_2|$ for BPF). For instance, take a density operator ρ described by a correlation vector $\vec{c}$ such that $|c_1| > |c_2| > |c_3| \neq 0$. By passing ρ through the PF channel, we will initially have $D_G[\varepsilon(\rho)] = |c'_2| = |c_2|(1-p)^2$. Then, there will be a first crossing involving $|c'_2|$ and the constant component $|c'_3| = |c_3|$ for a critical point $p = p_1^{SC}$, which will imply $D_G[\varepsilon(\rho)] = |c_3|$. The 1-norm GQD will be *frozen* as a function of p for a certain interval (robust to decoherence in this interval) and finally a new crossing will occur involving now $|c'_3|$ and $|c'_1|$ for a second critical point $p = p_2^{SC}$. From this point on, $D_G[\varepsilon(\rho)] = |c'_1| = |c_1|(1-p)^2$ until $p = 1$. Similar examples can be obtained for BF and BPF by replacing c_3 for c_1 and c_2, respectively. We also observe here that, in presence of symmetry, the double sudden change phenomenon may get degenerate into a single sudden change. As a simple example, this will happen in case of $U(1)$-symmetric states, for which $c_1 = c_2$. By passing such a state through the PF channel, we will have at most a single sudden change (if $|c_3| < |c_1| = |c_2|$).

(ii) GAD: Double sudden changes occur if and only if $|c_3| > |c_1|, |c_2|$ and $|c_1| \neq |c_2| \neq 0$. In this case, by passing ρ through the GAD channel, we will initially have $D_G[\varepsilon(\rho)] = \text{Max}[|c'_1|, |c'_2|] = \text{Max}[|c_1|, |c_2|](1-\gamma)$. Then, there will be a first crossing involving $\text{Max}[|c'_1|, |c'_2|]$ and the component $|c'_3|$ for a critical point $\gamma = p_1^{SC}$, which will imply $D_G[\varepsilon(\rho)] = |c'_3| = |c_3|(1-\gamma)^2$. Finally, a new crossing will occur involving now $|c'_3|$ and $\text{Min}[|c'_1|, |c'_2|]$ for a second critical point $\gamma = p_2^{SC}$. From this point on, $D_G[\varepsilon(\rho)] = \text{Min}[|c'_1|, |c'_2|] = \text{Min}[|c_1|, |c_2|](1-\gamma)$ until $\gamma = 1$. Note that no freezing occurs here. We also observe that $U(1)$-symmetric states also constrain the dynamics to exhibit at most a single sudden change (if $|c_3| > |c_1| = |c_2|$).

Hence, the 1-norm GQD exhibits two new aspects for nontrivial Bell-diagonal states $[|c_i| \neq 0\ (\forall i)]$ under Markovian evolution in comparison with the 2-norm GQD: (a) double sudden change as a function of its decay rate; (b) freezing behavior – and therefore robustness against decoherence – for a finite interval as a function of p for the channels BF, PF, and BPF. These properties can be obtained for the 2-norm GQD only if either more complex states are considered (e.g., X states) or the system is subjected to non-Markovian dynamics [30–34]. Indeed, the impossibility of the 2-norm GQD to reveal double sudden change and freezing for Bell-diagonal states under the Markovian local channels considered in this work can be proven below.

Proposition 1. *Consider a nontrivial two-qubit Bell-diagonal state $[\ |c_i| \neq 0\ (\forall i)\ ]$ evolving under Bell-diagonal-preserving Markovian local noise described by the BF, BPF, PF, or GAD channels. For such individual dynamical evolutions, the 2-norm GQD is unable to exhibit either double sudden change or nonvanishing freezing as a function of the decoherence parameters.*

Proof. From Ref. [24], we can write the 2-norm GQD as $D_2 = (1/4)(\sum_i |c'_i|^2 - \max |c'_i|^2)$. Then, in order for a double sudden change to occur, we must have two changes in the maximum value of $|c'_i|^2$ as a function of the decoherence parameter, namely, p for BF, PBF, PF, and γ for GAD. This is because $\sum_i |c'_i|^2$ is a smooth function of p or γ. However, from Table II, it can be observed that, for any channel, two of the correlation functions $|c'_i|^2$ display the same decay rate, which means that they do not cross as a function of p or γ. Therefore, at most one sudden change is allowed for the 2-norm GQD. Concerning the freezing behavior, we must have D_2 as a constant for a fixed interval of p or γ. By rewriting D_2 as $D_2 = (1/4)(\min |c'_i|^2 + \text{int}|c'_i|^2)$, it follows from Table II that D_2 cannot be a constant. Indeed, as discussed before, two of the correlation functions c'_i depend on p with the same decay rate. Moreover, the possibility of two nonvanishing constant values for c'_i has been excluded. Hence, there are no double sudden changes or freezing of correlations for the 2-norm GQD in the case of nontrivial Bell-diagonal states $[|c_i| \neq 0\ (\forall i)]$ and for the dynamical evolutions in Table II. □

Notice that, by relaxing the requirement $|c_i| \neq 0$ $(\forall i)$, we can obtain the freezing behavior for the 2-norm GQD D_2 for *separable* Bell-diagonal states (see, e.g., Ref. [35]). In particular, from $D_2 = (1/4)(\min|c_i'|^2 + \mathrm{int}|c_i'|^2)$, the freezing would occur for the BF, PF, and PBF channels by taking the constant component $|c_k|$ as the intermediate absolute value of the components of $\vec{c}$ and $\min|c_i| = 0$ $(i \neq k)$. However, this is a very restrictive solution, which is monotonically related to the 1-norm GQD, i.e., $D_2 = D_G^2/4$. In other words, the 1-norm GQD allows for the freezing behavior in a much wider class of Bell-diagonal states. A similar freezing behavior is also known for the entropic QD, where an interpretation of sudden transition between classical and quantum decoherence has been introduced [36]. The sudden change critical points $p = p_1^{SC}$ and $p = p_2^{SC}$ for the 1-norm GQD can be analytically obtained by imposing the equality of the absolute values of the components of $\vec{c'}$ at the crossings described in the discussion above. The results are organized in Table III.

	p_1^{SC}	p_2^{SC}	Conditions
BF	$1-\sqrt{\frac{\lvert c_1\rvert}{\mathrm{Min}(\lvert c_2\rvert,\lvert c_3\rvert)}}$	$1-\sqrt{\frac{\lvert c_1\rvert}{\mathrm{Max}(\lvert c_2\rvert,\lvert c_3\rvert)}}$	$\lvert c_1\rvert < \lvert c_2\rvert, \lvert c_3\rvert$ $\lvert c_1\rvert \neq 0$ $\lvert c_2\rvert \neq \lvert c_3\rvert$
PF	$1-\sqrt{\frac{\lvert c_3\rvert}{\mathrm{Min}(\lvert c_1\rvert,\lvert c_2\rvert)}}$	$1-\sqrt{\frac{\lvert c_3\rvert}{\mathrm{Max}(\lvert c_1\rvert,\lvert c_2\rvert)}}$	$\lvert c_3\rvert < \lvert c_1\rvert, \lvert c_2\rvert$ $\lvert c_3\rvert \neq 0$ $\lvert c_1\rvert \neq \lvert c_2\rvert$
BPF	$1-\sqrt{\frac{\lvert c_2\rvert}{\mathrm{Min}(\lvert c_1\rvert,\lvert c_3\rvert)}}$	$1-\sqrt{\frac{\lvert c_2\rvert}{\mathrm{Max}(\lvert c_1\rvert,\lvert c_3\rvert)}}$	$\lvert c_2\rvert < \lvert c_1\rvert, \lvert c_3\rvert$ $\lvert c_2\rvert \neq 0$ $\lvert c_1\rvert \neq \lvert c_3\rvert$
GAD	$1-\frac{\mathrm{Max}(\lvert c_1\rvert,\lvert c_2\rvert)}{\lvert c_3\rvert}$	$1-\frac{\mathrm{Min}(\lvert c_1\rvert,\lvert c_2\rvert)}{\lvert c_3\rvert}$	$\lvert c_3\rvert > \lvert c_1\rvert, \lvert c_2\rvert$ $\lvert c_1\rvert, \lvert c_2\rvert \neq 0$ $\lvert c_1\rvert \neq \lvert c_2\rvert$

TABLE III. Critical points p_1^{SC} and p_2^{SC} in terms of the components of the correlation vector $\vec{c}$. The conditions provided are necessary and sufficient for assuring double sudden changes. Notice that double sudden changes may get degenerate if symmetries (implying in the equality of some of the components of $\vec{c}$) are present in the quantum state.

IV. EXAMPLES

A. Double sudden change for the GAD channel

We begin by illustrating the double sudden change behavior for the GAD channel. Let us take a quantum system described by a Bell-diagonal state ρ. In order for ρ to correspond to a physical state, we must impose $\lambda_i \geq 0$ and $\sum_i \lambda_i = 1$, with λ_i denoting the eigenvalues of ρ. Under these conditions, the vector $\vec{c}$ turns out to be restricted to a tetrahedron whose vertices situated on the points $(1,1,-1)$, $(-1,-1,-1)$, $(1,-1,1)$, and $(-1,1,1)$, as plotted in Fig. 1. Then, by using the conditions obtained in Table III for the GAD channel, we can determine a physical region in the state space associated with double sudden changes in the decay rates of the 1-norm GQD. This is plotted in red color in Fig. 1. Within this physical region, we identify an explicit example to exhibit the double sudden change phenomenon, which is given by the values $c_1 = 0.1$, $c_2 = 0.2$, and $c_3 = 0.3$ (blue dot in Fig. 1). Its 1-norm GQD is then plotted in Fig. 2. From this plot, we observe the two critical points at $p_1^{SC} = 1/3$ and $p_2^{SC} = 2/3$, which are in agreement with the values in Table III.



FIG. 1. (Color online) Bell-diagonal states exhibiting double sudden changes under the GAD channel (red region inside the tetrahedron). The blue point represents the state $\vec{c} = (0.1, 0.2, 0.3)$ used as the initial state in Fig. 2.

B. Freezing behavior for the PF channel

Let us now illustrate the freezing behavior by using the PF channel. In this direction, we choose the state $c_1 = 1$ and $c_2 = -c_3$ (with $c_3 \neq 0$), which has been discussed in Ref. [36] for the entropic QD. For this state, it is possible to show that the freezing behavior is followed by a sudden change at $p^{SC} = 1 - \sqrt{|c_3|}$ for both the entropic QD and the 1-norm GQD. Notice that the 2-norm GQD does not exhibit freezing for this state, since $|c_i| \neq 0$ $(\forall i)$. Then, for the entropic QD and the 1-norm GQD, we obtain a large and controllable freezing interval. Indeed, by reducing $|c_3|$ (and consequently D_G), we can increase the freezing interval as close as desired to $p = 1$. Naturally, the longer the freezing interval is, the less 1-norm GQD is available. This is illustrated in Fig. 3, where we have chosen $c_3 = 0.1$. We observe that a large freezing is obtained, with the sudden change occurring for $p > 0.6$.



FIG. 2. (Color online) Double sudden change for the state given by $\vec{c}=(0.1,0.2,0.3)$ under the GAD channel, with $p_1^{SC}=1/3$ and $p_2^{SC}=2/3$. Inset: Absolute value of the correlation functions as a function of γ.



FIG. 3. (Color online) Freezing behavior for the state given by $\vec{c}=(1,-0.1,0.1)$ under the PF channel. Inset: Absolute value of the correlation functions as a function of p.

C. Quantum spin chains under decoherence

We consider now the correlation properties of a multi-qubit system under decoherence. More specifically, we will take a critical quantum spin chain, whose ground state is described by a Bell-diagonal state. In this context, we will illustrate the discussion of the 1-norm GQD between two spins by investigating the XXZ spin chain, whose Hamiltonian is given by

$$H_{XXZ} = -\frac{J}{2}\sum_{i=1}^{L}\left(\sigma_i^x\sigma_{i+1}^x + \sigma_i^y\sigma_{i+1}^y + \Delta\sigma_i^z\sigma_{i+1}^z\right), \quad (11)$$

where periodic boundary conditions are assumed, ensuring therefore translation symmetry. We will set the energy scale such that $J=1$ and will be interested in a nearest-neighbor spin pair at sites i and $i+1$. The model exhibits a first-order quantum phase transition at $\Delta=1$ and an infinite-order quantum phase transition at $\Delta=-1$. The entropic quantum discord has already been analyzed for nearest-neighbors, with the characterization of its quantum critical properties discussed [37, 38]. For the geometric versions of QD in the XXZ model, see Ref. [21]. Concerning its symmetries, the XXZ chain exhibits $U(1)$ invariance, namely, $\left[H,\sum_i\sigma_z^i\right]=0$, which ensures that the two-spin density operator, for the chain in its ground state, is given by Eq. (4). By using the Hellmann-Feynman theorem [39, 40] for the XXZ Hamiltonian (11), we then obtain

$$\begin{aligned} c_1 = c_2 &= \frac{1}{2}\left(G_{xx}+G_{yy}\right) = \Delta\frac{\partial\varepsilon_{xxz}}{\partial\Delta} - \varepsilon_{xxz}\,, \\ c_3 &= G_{zz} = -2\frac{\partial\varepsilon_{xxz}}{\partial\Delta}\,, \end{aligned} \quad (12)$$

where ε_{xxz} is the ground state energy density

$$\varepsilon_{xxz} = \frac{\langle\psi_0|H_{XXZ}|\psi_0\rangle}{L} = -\frac{1}{2}\left(G_{xx}+G_{yy}+\Delta G_{zz}\right), \quad (13)$$

with $|\psi_0\rangle$ denoting the ground state of H_{XXZ}. Eqs. (12) and (13) hold for a chain with an arbitrary number of sites, allowing the discussion of correlations either for finite or infinite chains. Indeed, ground state energy as well as its derivatives can be exactly determined by Bethe Ansatz technique for a chain in the thermodynamic limit [41]. Then, from Tables II and III, we can obtain a general picture of the decoherence effects for different quantum phases. In particular, it is possible to show (see, e.g., Ref. [37]) that $c_1 < c_3$ for both $\Delta < -1$ and $\Delta > 1$. On the other hand, $c_1 > c_3$ for the gapless disordered phase $-1 < \Delta < 1$. This implies, for the XXZ model, that the occurrence of sudden changes may be driven by quantum phase transitions, whose behavior is summarized in Table IV.

Channel	$\Delta < -1$	$-1 < \Delta < 1$	$\Delta > 1$
BF	1 S.C.	No S.C.	No S.C.
PF	No S.C.	No S.C.	No S.C.
BPF	1 S.C.	No S.C.	No S.C.
GAD	1 S.C.	No S.C.	No S.C.

TABLE IV. Occurrence of sudden changes (denoted here as S.C.) in the 1-norm GQD for different quantum channels. The decoherence behavior may change for different quantum phases. The phase $\Delta > 1$ does not exhibit S.C. because $c_1 = c_2 = 0$.

For a specific example, we consider the decoherence behavior for the BF flip channel in the three quantum regimes governed by the anisotropy Δ. This is plotted in Fig. 4. As we can see, the sudden change behavior that is present in the antiferromagnetic phase ($\Delta = -1.5$ in the plot) disappears in the gapless phase ($\Delta = 0$ in the plot), characterizing the infinite-order quantum phase transition of the XXZ chain. Therefore, decoherence turns out to be a probe for the antiferromagnetic quantum critical point. Such a characterization is a particular feature of the XXZ chain, holding possibly also for related models, but not implying a general approach for quantum criticality. For the entropic QD, a detection of quantum phase transitions in terms of sudden change behavior has recently been introduced in Refs. [42, 43].



FIG. 4. (Color online) Sudden change behavior for first neighbors in the ground state of the XXZ chain under the BF channel. Distinct phases may exhibit different behaviors, going from monotonic decay ($\Delta > -1$) to sudden change dynamics ($\Delta < -1$).

V. CONCLUSIONS

We have analytically investigated the 1-norm GQD under the effect of Markovian noise for Bell-diagonal states. In particular, we have shown that, for the simplest cases of Markovian dynamics, the 1-norm GQD may already exhibit double sudden changes or freezing behavior. The origin of such features is the robustness of QD under decoherence. This inherent robustness can be traced back as a consequence of the negligibility of the set of classical states (zero discord states) in comparison with the whole state space [47]. This is in strong contrast with the set of separable (nonentangled) states, which exhibits finite volume in state space. This implies, e.g. in the possibility of sudden death for entanglement, while just sudden changes (typically with no death) in the case of QD. Specifically concerning the 1-norm GQD, it inherits this robustness of QD against noise and reveals it for simple Bell-diagonal states, which is a by-product of the trace norm properties. Experimental investigations of single sudden changes and freezing behavior have been realized through different physical setups [44–46], which have demonstrated that such phenomena are quite general in real conditions. Remarkably, a recent investigation [28] experimentally verified the single sudden change and the correlation freezing for the negativity of quantumness in Bell-diagonal states, which turns out to be equivalent to the 1-norm GQD. Therefore, further experiments aiming at those properties and also at double sudden changes could greatly benefit from the correlation dynamics derived in this work. Moreover, we have found a surprising connection between the sudden change behavior and the infinite-order quantum phase transition in the XXZ model. Such kind of connections appear as a promising direction for investigating quantum criticality. We leave these possibilities as future challenges, as well as generalizations for more general states and for non-Markovian dynamics.

ACKNOWLEDGMENTS

This work is supported by the Brazilian agencies CNPq, CAPES, FAPERJ, and the Brazilian National Institute for Science and Technology of Quantum Information (INCT-IQ).

---

[1] V. Vedral, Rev. Mod. Phys. **74**, 197 (2002).
[2] A. Shimony, Ann. NY. Acad. Sci. **755**, 675 (1995).
[3] H. Barnum and N. Linden, J. Phys. A: Math. Gen. **34**, 6787 (2001).
[4] T.-C. Wei, P. M. Goldbart, Phys. Rev. A **68**, 042307 (2003).
[5] K. Modi, A. Brodutch, H. Cable, T. Paterek, V. Vedral, Rev. Mod. Phys. **84**, 1655 (2012).
[6] L. C. Céleri, J. Maziero, R. M. Serra, Int. J. Quantum Inf. **9**, 1837 (2011).
[7] M. S. Sarandy, T. R. de Oliveira, L. Amico, Int. J. Mod. Phys. B **27**, 1345030 (2013).
[8] H. Ollivier, W. Zurek, Phys. Rev. Lett. **88**, 017901 (2001).
[9] B. Dakić, V. Vedral, C. Brukner, Phys. Rev. Lett. **105**, 190502 (2010).

[10] A. C. S. Costa, R. M. Angelo, Phys. Rev. A **87**, 032109 (2013).
[11] S. Luo, S. Fu, Phys. Rev. A **82**, 034302 (2010).
[12] A. S. M. Hassan, B. Lari, P. S. Joag, Phys. Rev. A **85**, 024302 (2012).
[13] S. Rana, P. Parashar, Phys. Rev. A **85**, 024102 (2012).
[14] B. Dakic, Y. O. Lipp, X. Ma, M. Ringbauer, S. Kropatschek, S. Barz, T. Paterek, V. Vedral, A. Zeilinger, C. Brukner, P. Walther, Nature Phys. **8**, 666 (2012).
[15] M. Gu, H. M. Chrzanowski, S. M. Assad, T. Symul, K. Modi, T. C. Ralph, V. Vedral, P. K. Lam, Nature Phys. **8**, 671 (2012).
[16] B. Bellomo, G. L. Giorgi, F. Galve, R. Lo Franco, G. Compagno, R. Zambrini, Phys. Rev. A **85**, 032104 (2012).
[17] X. Hu, H. Fan, D. L. Zhou, W.-M. Liu, arXiv:1203.6149 (2012).
[18] T. Tufarelli, D. Girolami, R. Vasile, S. Bose, G. Adesso, Phys. Rev. A **86**, 052326 (2012).
[19] M. Piani, Phys. Rev. A **86**, 034101 (2012).
[20] T. Debarba, T. O. Maciel, R. O. Vianna, Phys. Rev. A **86**,024302 (2012).
[21] F. M. Paula, T. R. de Oliveira, M. S. Sarandy, arXiv:1302.7034 (2013).
[22] S. Rana, P. Parashar, Phys. Rev. A **87**, 016301 (2013).
[23] J. Maziero, L. C. Céleri, R. M. Serra, V. Vedral, Phys. Rev. A **80**, 044102 (2009).
[24] X.-M. Lu, Z.-J. Xi, Z. Sun, X. Wang, Quantum Inform. Comput. **10**, 994 (2010).
[25] K. Modi, T. Paterek, W. Son, V. Vedral, M. Williamson, Phys. Rev. Lett. **104**, 080501 (2010).
[26] M. Piani, S. Gharibian, G. Adesso, J. Calsamiglia, P. Horodecki, A. Winter, Phys. Rev. Lett. **106**, 220403 (2011).
[27] T. Nakano, M. Piani, G. Adesso, arXiv:1211.4022 (2012).
[28] I. A. Silva, D. Girolami, R. Auccaise, R. S. Sarthour, I. S. Oliveira, T. J. Bonagamba, E. R. deAzevedo, D. O. Soares-Pinto, G. Adesso, Phys. Rev. Lett **110**, 140501 (2013).
[29] M. A. Nielsen, I. L. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, 2000.
[30] G. Karpat, Z. Gedik, Phys. Lett. A **375**, 4166 (2011).
[31] W. Song, L.-B. Yu, P. Dong, D.-C. Li, M. Yang, Z.-L. Cao, arXiv:1112.4318 (2011).
[32] W. Song, L.-B. Yu, D.-C. Li, P. Dong, M. Yang, Z.-L. Cao, arXiv:1203.3356 (2012).
[33] H. Yuan, L.-F. Wei, Int. J. Theor. Phys. **52**, 987 (2013).
[34] R. Lo Franco, B. Bellomo, S. Maniscalco, G. Compagno, Int. J. Mod. Phys. B **27**, 1345053 (2013).
[35] Y. Yao, H.-W. Li, Z.-Q. Yin, Z.-F. Han, Phys. Lett. A **376**, 358 (2012).
[36] L. Mazzola, J. Piilo, S. Maniscalco, Phys. Rev. Lett. **104**, 200401 (2010).
[37] M. S. Sarandy, Phys. Rev. A **80**, 022108 (2009).
[38] R. Dillenschneider, Phys. Rev. B **78**, 224413 (2008).
[39] H. Hellmann, *Die Einführung in die Quantenchemie* (Deuticke, Leipzig, 1937).
[40] R. P. Feynman, Phys. Rev. **56**, 340 (1939).
[41] C. N. Yang, C. P. Yang, Phys. Rev. **150**, 321 (1966); *ibid.* **150**, 327 (1966).
[42] A. K. Pal, I. Bose, Eur. Phys. J. B **85**, 277 (2012).
[43] I. Bose, A. K. Pal, Int. J. Mod. Phys. B **27**, 1345042 (2013).
[44] J.-S. Xu, C.-F. Li, C.-J. Zhang, X.-Y. Xu, Y.-S. Zhang, G-C. Guo, Phys. Rev. A **82**, 042328 (2010).
[45] J.-S. Xu, X.-Y. Xu, C.-F. Li, C.-J. Zhang, X.-B. Zou, G.-C. Guo, Nature Commun. **1**, 7 (2010).
[46] R. Auccaise, L. C. Céleri, D. O. Soares-Pinto, E. R. deAzevedo, J. Maziero, A. M. Souza, T. J. Bonagamba, R. S. Sarthour, I. S. Oliveira, R. M. Serra, Phys. Rev. Lett. **107**, 140403 (2011).
[47] A. Ferraro, L. Aolita, D. Cavalcanti, F. M. Cucchietti, A. Acín, Phys. Rev. A **81**, 052318 (2010).